

DAFTAR PUSTAKA

- [1] American Cancer Society, “Key Statistics About Kidney Cancer,” <https://www.cancer.org/cancer/types/kidney-cancer/about/key-statistics.html>, 2024, diakses 22 Oktober 2025.
- [2] Parkway Cancer Centre Indonesia, “Perkembangan Baru dalam Pengobatan Kanker Ginjal,” <https://www.parkwaycancercentre.com/id/idn/news-events/news-articles/news-articles-details/perkembangan-baru-dalam-pengobatan-kanker-ginjal>, 2022, diakses 23 Desember 2025.
- [3] H. N. Cahyanto, O. Zulkarnain, and R. Rahagia, “Pengembangan deteksi dini dan asuhan keperawatan pada kanker menggunakan artificial intelligence (ai) berbasis web,” *PREPOTIF Jurnal Kesehatan Masyarakat*, vol. 8, no. 3, pp. 7511–7518, December 2024, lisensi CC BY-SA 4.0. [Online]. Available: <https://doi.org/10.31004/prepotif.v8i3.34374>
- [4] A. Dewantoro, W. C. Anggundari, U. Nuraeni, B. Prasetya *et al.*, “Metode deteksi molekuler berbasis genomik dan diagnostik akurasi dalam pengembangan diagnostik klinik di indonesia,” in *Pertemuan dan Presentasi Ilmiah Standardisasi 2020*. Badan Riset dan Inovasi Nasional, February 2021, pp. 207–216. [Online]. Available: <https://doi.org/10.31153/ppis.2020.92>
- [5] Masyarakat Bioinformatika dan Biodiversitas Indonesia, “Bioinformatika Keanekaragaman Hayati: Menautkan Ekosistem, Genom, dan Kesehatan Manusia,” <https://mabbi.id/2025/11/03/bioinformatika-keanekaragaman-hayati-menautkan-ekosistem-genom-dan-kesehatan-manusia/> November 3 2025, diakses 23 Desember 2025.
- [6] Columbia Asia Indonesia, “Tren Terkini dalam Pengobatan Kanker: Terapi Inovatif dan Penelitian Baru,” <https://columbiaasia.co.id/artikel/kesehatan/tren-dalam-pengobatan-kanker/>, September 24 2024, diakses 23 Desember 2025.
- [7] National Cancer Institute (NCI) and National Human Genome Research Institute (NHGRI), “The Cancer Genome Atlas (TCGA) Homepage,” <https://cancergenome.nih.gov/>, 2009, diakses 28 April 2009.
- [8] A. M. Ali, H. Zhuang, A. Ibrahim, O. Rehman, M. Huang, and A. Wu, “A machine learning approach for the classification of kidney cancer subtypes using mirna genome data,” *Sensors*, vol. 18, no. 11, pp. 1–15, November 2018, published 29 November 2018.

- [9] Faj Lennon, “Memahami Lasso: Panduan Lengkap untuk Pemula,” <https://streamanalysis.com/blog/memahami-lasso-panduan-lengkap-untuk>, October 23 2025, diakses 23 Desember 2025.
- [10] S. Hochreiter and J. Schmidhuber, “Long short-term memory,” *Neural Computation*, vol. 9, no. 8, pp. 1735–1780, 1997. [Online]. Available: <https://doi.org/10.1162/neco.1997.9.8.1735>
- [11] A. K. Ibrahim, H. Zhuang, L. M. Chérubin, M. T. Schärer-Umpierre, and N. Erdol, “Automatic classification of grouper species by their sounds using deep neural networks,” *The Journal of the Acoustical Society of America*, vol. 144, no. 3, pp. EL196–EL202, 2018. [Online]. Available: <https://doi.org/10.1121/1.5054916>
- [12] B. W. Matthews, “Comparison of the predicted and observed secondary structure of t4 phage lysozyme,” *Biochimica et Biophysica Acta (BBA) - Protein Structure*, vol. 405, no. 2, pp. 442–451, 1975. [Online]. Available: [https://doi.org/10.1016/0005-2795\(75\)90109-9](https://doi.org/10.1016/0005-2795(75)90109-9)
- [13] R. L. Siegel, T. B. Kratzer, A. N. Giaquinto, H. Sung, and A. Jemal, “Cancer statistics, 2025,” *CA: A Cancer Journal for Clinicians*, 2025. [Online]. Available: <https://doi.org/10.3322/caac.21871>
- [14] M. Daugherty, S. Blakely, O. Shapiro, S. Vourganti, M. Mollapour, and G. Bratslavsky, “Chromophobe renal cell carcinoma is the most common nonclear renal cell carcinoma in young women: results from the seer database,” *Journal of Urology*, vol. 195, no. 4 Part 1, pp. 847–851, 2016. [Online]. Available: <https://doi.org/10.1016/j.juro.2016.05.041>
- [15] American Cancer Society (ACS), “Kidney cancer staging,” <https://www.cancer.org/cancer/types/kidney-cancer/detection-diagnosis-staging/staging.html>, 2024, diakses 22 Oktober 2025.
- [16] Cancer Research UK, “The number stages of kidney cancer,” <https://www.cancerresearchuk.org/about-cancer/kidney-cancer/stages-types-grades/number-stages>, 2024, diakses 22 Oktober 2025.
- [17] R. C. Friedman, K. K.-H. Farh, C. B. Burge, and D. P. Bartel, “Most mammalian mRNAs are conserved targets of microRNAs,” *Genome Research*, vol. 19, no. 1, pp. 92–105, 2009. [Online]. Available: <https://doi.org/10.1101/gr.082701.108>
- [18] V. Ambros, “The functions of animal microRNAs,” *Nature*, vol. 431, no. 7006, pp. 350–355, 2004. [Online]. Available: <https://doi.org/10.1038/nature02871>
- [19] M. D. Jansson and A. H. Lund, “MicroRNA and cancer,” *Molecular Oncology*, vol. 6, no. 6, pp. 590–610, 2012. [Online]. Available: <https://doi.org/10.1016/j.molonc.2012.09.006>

- [20] S. Volinia, G. A. Calin, C.-G. Liu, S. Ambs, A. Cimmino, F. Petrocca, R. Visone, M. Iorio, C. Roldo, M. Ferracin *et al.*, “A microRNA expression signature of human solid tumors defines cancer gene targets,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 103, no. 7, pp. 2257–2261, 2006. [Online]. Available: <https://doi.org/10.1073/pnas.0510565103>
- [21] C. M. Croce, “Causes and consequences of microRNA dysregulation in cancer,” *Nature Reviews Genetics*, vol. 10, no. 10, pp. 704–714, 2009. [Online]. Available: <https://doi.org/10.1038/nrg2634>
- [22] E. I. Papadopoulos, C. Petraki, A. Gregorakis, E. G. Fragoulis, and A. Scorilas, “Clinical evaluation of microRNA-145 expression in renal cell carcinoma: A promising molecular marker for discriminating and staging the clear cell histological subtype,” *Biological Chemistry*, vol. 397, no. 6, pp. 529–539, 2016. [Online]. Available: <https://doi.org/10.1515/hsz-2015-0296>
- [23] N. M. White, T. T. Bao, J. Grigull, Y. M. Youssef, A. Girgis, M. Diamandis, E. Fatoohi, M. Metias, R. J. Honey, R. Stewart *et al.*, “miRNA profiling for clear cell renal cell carcinoma: Biomarker discovery and identification of potential controls and consequences of miRNA dysregulation,” *The Journal of Urology*, vol. 186, no. 3, pp. 1077–1083, 2011. [Online]. Available: <https://doi.org/10.1016/j.juro.2011.04.044>
- [24] D. Juan, G. Alexe, T. Antes, H. Liu, A. Madabhushi, C. Delisi, S. Ganesan, G. Bhanot, and L.-S. Liou, “Identification of a microRNA panel for clear-cell kidney cancer,” *Urology*, vol. 75, no. 4, pp. 835–841, 2010. [Online]. Available: <https://doi.org/10.1016/j.urology.2009.09.085>
- [25] S. Samaan, H. W. Khella, A. Girgis, A. Scorilas, E. Lianidou, M. Gabril, S. N. Krylov, M. Jewett, G. A. Bjarnason, H. El-Said *et al.*, “mir-210 is a prognostic marker in clear cell renal cell carcinoma,” *The Journal of Molecular Diagnostics*, vol. 17, no. 2, pp. 136–144, 2015. [Online]. Available: <https://doi.org/10.1016/j.jmoldx.2014.10.003>
- [26] M. Desai and M. Shah, “An anatomization on breast cancer detection and diagnosis employing multi-layer perceptron neural network (mlp) and convolutional neural network (cnn),” *Clinical eHealth*, vol. 4, pp. 1–11, 2021.
- [27] H. Ramchoun, M. Amine, J. Idrissi, Y. Ghanou, and M. Ettaouil, “Multilayer perceptron: Architecture optimization and training,” *International Journal of Interactive Multimedia and Artificial Intelligence*, vol. 4, no. 1, p. 26, 2016.